

Figure 1

The Enterohepatic Circulation with Key Transporter Proteins Mediating

Bile Acid Circulation





Figure 2

Bile Acid Prodrug Derivatives for Sustained Release of Drugs



Y^a , Y^b are cleavable linker groups

D is a drug moiety

Q is CH_2 or O

W is selected from the group consisting of $-CH(CH_3)W'$ where W' is a substituted alkyl group containing a moiety which is negatively charged at physiological pH, which moiety is selected from the group consisting of $-COOH$, $-SO_3H$, $-SO_2H$, $-P(O)(OR^6)(OH)$, $-OP(O)(OR^6)(OH)$, $-OSO_3H$ and pharmaceutically acceptable salts thereof.

$R1 = R2 = \alpha-OH$ (from Cholate)

$R1 = \alpha-OH$, $R2 = H$ (from Chenodeoxycholate)

$R1 = \beta-OH$, $R2 = H$ (from Ursodeoxycholate)


$R1 = H$, $R2 = \alpha-OH$ (from Deoxycholate)

$R1 = \beta-OH$, $R2 = \alpha-OH$ (from Ursocholate)


$R1 = R2 = H$ (from Lithocholate)

Figure 3- Generic Structures of Preferred Bile Acid C-3 Derivatives

Hydroxyl or 1° and 2° Amine-Containing Drugs





Carboxylic Acid-Containing Drugs



W^{'''} is OH, NHCH₂CO₂H, NHCH₂CH₂SO₃H or pharmaceutically acceptable salts thereof

Figure 4- Generic Structures of Preferred Bile Acid C-24 Derivatives

Hydroxyl or 1° and 2° Amine-Containing Drugs



Q = O, CH₂; M = O, NR⁷

Carboxylic Acid-Containing Drugs





Figure 5 GABA Analog Derivatives and L-Dopa Derivatives



Generalized GABA Analog

Optionally Protected L-Dopa Analog

R14, R15, R16, R19 and R20 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl;


R17 and R18 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, acyl, substituted acyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl or optionally, R17 and R18 together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocycloalkyl, substituted heterocycloalkyl or bridged cycloalkyl ring;

P is a catechol protecting group (see Figure 6)

The GABA analog or L-Dopa analog is attached to the steroid nucleus in (I-a) or (I-b) either by replacement of one of the amino hydrogen atoms, or a hydrogen atom from one of the hydroxy groups of the catechol, or the hydroxyl group of the carboxyl moiety by a covalent bond to Y^a or Y^b



Figure 6:

Catechol Protection Strategies Applicable for L-Dopa Bile Acid Conjugates



R30 = hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl
 R31 = alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl
 R24, R25 = hydrogen, alkyl, substituted alkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl
 or R24 and R25 together with the carbon to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocycloalkyl or substituted heterocycloalkyl ring

1. *Chrysomelidae* (10 species)
 2. *Curculionidae* (10 species)
 3. *Chrysomelidae* (10 species)
 4. *Curculionidae* (10 species)
 5. *Chrysomelidae* (10 species)
 6. *Curculionidae* (10 species)
 7. *Chrysomelidae* (10 species)
 8. *Curculionidae* (10 species)
 9. *Chrysomelidae* (10 species)
 10. *Curculionidae* (10 species)



m' is 0 to 6 ; n' is 0 to 6

K = 0, NR7, CR8R9; S(0)_j, j = 0, 1, or 2

R26 and R27 are independently selected from the group consisting of halo and lower alkyl (including branched alkyl)

Figure 8
Enterohepatic Circulation Mediated by Intestinal Peptide
and Hepatic Anion Transporters





Figure 9

Enterohepatic Recirculating Prodrugs Based On Glutathione Mimetics



R13 = H , lower alkyl

Use PEPT1 substrate with metabolically stable di- or tripeptide backbone to achieve intestinal absorption

Figure 10





Figure 11



(Faint, illegible handwritten notes)




Figure 13





Figure 14 - Synthesis of Cholyl-Dopa Conjugates



1. The first of these is the fact that the
 2. of the first of these is the fact that the
 3. of the first of these is the fact that the
 4. of the first of these is the fact that the
 5. of the first of these is the fact that the
 6. of the first of these is the fact that the
 7. of the first of these is the fact that the
 8. of the first of these is the fact that the
 9. of the first of these is the fact that the
 10. of the first of these is the fact that the












Figure 16



[illegible]